

Larry Crouse

Extension Horticulture Agent

PEPPERS FROM THE GARDEN

Peppers are able to be stored fresh much longer than tomatoes. They can usually keep in a crisper drawer of a refrigerator for several weeks if kept moist but not wet. For longer storage, freezing works well. Though mushy when thawed, the flavor still comes through in cooked foods. Try dicing them into small pieces and then freezing on a cookie sheet. The frozen pieces can then be poured into a plastic bag for later use. Measuring is much easier as the pieces are not frozen together in a clump. This method works equally well for hot peppers but be sure to wear gloves when handling.

Should You Let Turf Grow Tall in the Fall?

Sometimes you will hear people say to let the grass grow tall right before winter sets in. Their reasoning is that the extra foliage will insulate the crown of the plant from the extreme cold of winter. Although this may sound reasonable, in practice it probably does little, if anything, to increase winter hardiness. On the contrary, a canopy that is too high during the winter may lay over and become matted down, leading to an increased incidence of winter-diseases such as snow mold. Turfgrass species vary genetically in their cold tolerance, with warm-season grasses such as bermudagrass, zoysiagrass and buffalograss being less cold tolerant than the cool-season types such as tall fescue and Kentucky bluegrass. Given these differences, cold tolerance is improved by increasing the health of the plants going into the winter, and healthy plants are a result of a sound management program (fertilizing, watering and mowing) during the spring, summer and fall. The lawn will benefit more from continuing to mow at the recommended height than from trying to gain some insulation against winter cold by allowing it to grow tall. Here is a list of the recommended mowing height ranges (in inches) for home lawns in Kansas:

Tall fescue 2.5 -3.5; Kentucky bluegrass 2-3; Buffalograss 2-3; Bermudagrass 1-2; Zoysiagrass 1-2

(Note: Mowing at heights below 1.5 inches requires a reel mower).

There may be some benefits gained by adjusting mowing heights WITHIN the recommended range at times. For example, it is a good practice to mow warm-season grasses at the higher end of recommended heights during late summer and early fall because this practice should help them store more carbohydrate reserves for the winter, and it may reduce the incidence of certain cool-weather diseases. But the rule to remember is to stay within the recommended height range for your species.

Last Tomatoes of the Season

Cold nights are increasing in frequency now that we are into October. If you have tomatoes, you may have some that are approaching maturity. Leave them on the vine until mature or until a frost is forecast. Tomatoes will ripen off the vine but must have reached a certain phase of maturity called the 'mature green stage.' Look for full-sized tomatoes with a white, star-shaped zone on the bottom end of the green fruit. When harvesting fruit before a frost, separate tomatoes into three groups for storage: those that are mostly red, those that are just starting to turn, and those that are still green. Discard tomatoes with defects such as rots or breaks in the skin. Place the tomatoes on cardboard trays or cartons but use layers of newspaper to separate fruit if stacked. Occasionally a tomato may start to rot and leak juice. The newspaper will keep the juice from contacting nearby or underlying fruit. Store groups of tomatoes as close to 55 degrees as possible until needed.

Fall Planting of Asparagus & Rhubarb

We sometimes receive questions as to whether asparagus or rhubarb can be moved in the fall. Though these crops are traditionally transplanted in the spring (mid-March to mid-April), a fall move can be successful. Wait until the top has been browned by frost and then cut back to the ground. Prepare the soil and fertilize as you would in the spring.

See > <http://www.bookstore.ksre.ksu.edu/pubs/mf319.pdf>< for more detail on asparagus and > <http://www.bookstore.ksre.ksu.edu/pubs/ep99.pdf>< for more information on rhubarb.

Water well after planting to ensure good root/soil contact. Mulching would be helpful on the rhubarb to prevent the plant from heaving out of the soil during the winter but asparagus requires no such treatment as it is planted much deeper.

Mycosphaerella Leaf Spot on Ash

With all the concern about Emerald Ash Borer (EAB), many people may assume that any ash tree with problems is being attacked by EAB. However, EAB has only been confirmed in Atchison, Doniphan, Douglas, Jefferson, Johnson, Leavenworth, Shawnee & Wyandotte counties. One of the other problems we see with ash is *Mycosphaerella* Leafspot. Though this disease looks serious, it is not. *Mycosphaerella* leaf spot causes small, brown spots that enlarge to become blotches and may result in early leaf drop. This disease can be severe enough to cause complete defoliation of the tree. However, defoliation this late in the growing season will not hurt the health of the tree. Therefore, because this disease appears sporadically and tree health is not harmed, we do not recommend treatment. Furthermore, treatment would have to be preventative and applied before the disease had infected the leaves. Applying a fungicide now would have no effect.

Kansas State University Cooperative Extension Office

206 N. Griffith, Suite A

El Dorado, Kansas 67042

