

THE GRAPEVINE

November 10th, 2019

Larry Crouse

Extension Horticulture Agent

APPLY FERTILIZER APPLICATION IN NOVEMBER

November is the time to give Kentucky bluegrass and tall fescue lawns the last nitrogen application of the season. Why November? Because while top growth slows in response to cool temperatures, grass plants are still making food (carbohydrates) by photosynthesis. A November nitrogen application helps boost the photosynthesis rate. Carbohydrates that are not used in growth are stored in the crown and other storage tissues in the plant. These carbohydrate reserves help the turfgrass green up earlier in the spring and sustain growth into May without the need for early-spring (March or April) nitrogen. Those early-spring nitrogen applications are less desirable because they can lead to excessive shoot growth and reduced root growth. Other benefits of November-applied nitrogen for cool-season grasses include improved winter hardiness, root growth and shoot density. How much should you apply? One to 1 to 1 ½ pounds actual nitrogen per 1,000 sq. ft. of lawn area is sufficient. In order for this application to be effective the nitrogen must be readily available to the plant because the growing season is nearly over. Therefore, for a November application, use a soluble (quickly-available) nitrogen carrier such as urea or ammonium sulfate. Many turfgrass fertilizers sold in garden centers and other retail outlets also contain soluble nitrogen. Avoid products that contain water-insoluble nitrogen (slow-release) for this application. As always, sweep up any fertilizer that gets on driveways, sidewalks, or streets and reapply it to the lawn.

Garden Mums

As soon as garden chrysanthemums are done flowering, you may cut the plants back to 2 to 3 inches high. Some gardeners prefer to leave the top growth so that it provides some protection from fluctuating soil temperatures. If you choose to cut the tops off, apply a layer of mulch over the top of your mums after the ground has frozen or if the forecast calls for a sharp drop in temperature. Mums should not completely dry out during the winter. It may be necessary to water occasionally if sufficient rain or snow has not fallen.

What to Do with Tree Leaves

It's that time of year again. Leaves are rapidly falling from deciduous trees so it's a good time to stop and think about options for handling the litter. Although a scattering of leaves won't harm the lawn, excessive cover prevents sunlight from reaching turfgrass plants. Turf left in this state for an extended period will be unable to make the carbohydrates needed to carry it through the winter. There are options for dealing with the fallen leaves other than bagging them up and putting them out for the trash collector. Composting is a great way to handle the refuse. Compost can then be used in the vegetable garden and flowerbeds. An even easier method of making good use of the leaves is direct incorporation in either vegetable gardens or annual flower beds. Use a lawn mower with a bagging attachment to chop and collect the leaves. Transport them to the garden or bed and apply a 2 to 3-inch layer of leaves on the surface of the soil and then till them in. Repeat the process every couple of weeks until you run out of leaves or the weather becomes too cold or the soil becomes too wet. With luck, you should be able to make 3 to 4 applications this fall. Another option is to mow the leaves with a mulching mower and let shredded leaves filter into the turf canopy. (A side-discharge mower also will work, but it won't shred the

leaves as thoroughly.) This method will be most effective if you do it often enough that leaf litter doesn't become too thick. Mow while you can still see grass peeking through the leaves. You may wonder whether this practice will be detrimental to the lawn in the long run. Research at Michigan State University in which they used a mulching mower to shred up to about one pound of leaves per square yard of lawn (one pound is equal to approximately 6 inches of leaves piled on the grass) for five consecutive years, found no long-term effects of the shredded leaves on turf quality, thatch thickness, organic content of the thatch, or soil test results (pH, nutrients, etc.). If you mow leaves and have a cool-season lawn, it makes sense to be on a fall nitrogen fertilization program and core-aerate in the fall (things you should be doing anyway). If you have a warm-season lawn, you can still use this technique but wait to fertilize and core-aerate until next late May or early June

Horseradish

Horseradish is ready to dig after a hard freeze kills the foliage (usually November or December). The large roots can be harvested while smaller, pencil sized roots can be cut in 6-8 inch-long sections as 'seed' or 'sets' for next year's crop which are then immediately re-planted. Another option is to leave the horseradish in the ground and dig as needed. If you choose the latter option, be sure to heavily mulch the area so that the ground doesn't freeze. To use horseradish, peel the large, fleshy roots and cut into sections. Use a blender or food processor to chop the roots along with a small amount of water and a couple of ice cubes. Vinegar or lemon juice is added to stop the process that produces the "bite" of horseradish. Add immediately after blending for a mild flavor or wait up to 3 minutes to give the horseradish more kick. Use 2 to 3 tablespoons of vinegar or lemon juice per cup of horseradish sauce along with ½ teaspoon of salt for flavor. Horseradish has an extremely strong odor and so you may wish to open the blender or food processor outdoors and to keep your face away from the container when opening. Store ground horseradish in a tightly sealed jar in a refrigerator until ready for use.

Knotweed Control

Knotweed thrives in compacted soils, so a thorough aeration is the first step in control. This weed will not compete in a healthy lawn. Chemically, there are two options. Knotweed is an annual that germinates in late February or early March, so a preemergence herbicide can be used in the late fall (about now). Pendimethalin (Scotts Halts), Surflan (Weed Impede), Barricade, Dimension and XL are labeled for knotweed. (Note: Pendimethalin, Barricade and Dimension can be used on all Kansas turfgrasses, while Surflan and XL can only be used on tall fescue and warm-season grasses such as buffalograss, zoysiagrass and bermuda). The other option is to use a combination postemergence product such as Trimec, Weed-Out, Weed-B-Gon or Weed Free Zone after the knotweed has emerged in the spring but is still young. If spring seeding of the lawn is planned, your options are more limited. Buctril can be used on commercial sites and has a very short residual. It must be used on very young knotweed to get control. Trimec and others require a month before overseeding to thicken up your lawn. Obviously, don't use a preemergence herbicide if you are trying to get new seed established. For homeowners seeding in the spring, tilling will control knotweed adequately without using a herbicide.

Larry E. Crouse
Butler County Horticulture Agent
K-State Research & Extension
206 N Griffith, Suite A
El Dorado, KS 67042
(316)321-9660
lcrouse@ksu.edu