



January 9th, 2020 Larry Crouse **Extension Horticulture Agent**



STARTING ONION PLANTS INDOORS

It can be difficult to find specific onion varieties in sets or transplants, so growing from seed may be a preferred option. Onions are one of the first plants to be seeded for transplanting because this crop takes a significant amount of time (6 to 8 weeks) to reach transplant size and because they can be set out relatively early (late March in our area). Therefore, we want to start onions in mid- to late-January. Onion seed should be placed ½ to 3/4 inch apart in a pot or flat filled with a seed starting mix. Place the container in a warm (75 to 80 F) location until young seedlings emerge. Move to a cooler location (60 to 65 F) when the seedlings are 1 to 2 inches tall. Make sure they have plenty of light, using florescent or LED lights if needed. Start fertilizing when the seedlings reach 2 to 3 inches tall using a soluble fertilizer with each or every other watering. Onion seedlings tend to be spindly with the remains of the seed sticking to the end of a leaf for several weeks. Encourage stockiness by trimming the ends of the leaves when the plants reach 4 to 5 inches tall. Start hardening off the onions in early March by moving the plants to a protected outdoor location. Move them inside to protect them from extreme cold snaps.

Forcing Paperwhite Bulbs

Paperwhites are a form of daffodil that do not require a chilling period in order to bloom. Therefore, they are very easy to force and bring in to bloom. Following are the steps needed.

- o Use a 3 to 4-inch decorative container that does not have drainage holes. It should be transparent enough that you can see the water level in relation to the bulbs.
- o Place 1 to 2 inches of washed gravel, marbles, glass beads or stones in the bottom of the container. We will call the material chosen as "media" for the remainder of the article.
- o Place the bulbs on the media close to each other. Add enough media to hold them in place.
- o Add enough water that the bottom of the bulb is sitting in water. Do not submerge the bulb. Maintain the water at this level. It normally takes 4 to 8 weeks for the bulbs to bloom.

Unfortunately, paperwhites often become leggy and fall over. Growing in cooler temperatures (60 to 65 degrees) can help but there is another trick that can be useful and involves using a dilute solution of alcohol. No, this trick did not come from an unknown source on the Internet but Cornell University's Flower Bulb Research Program. They suggest the following to obtain a plant that is 1/3 shorter than normal. Flower size and longevity are not affected.


- o Grow the bulbs as described above until the shoot is green and about 1 to 2 inches above the top of the bulb.
- o Pour off the water and replace it with a 4 to 6% alcohol solution.
- o Use this solution instead of water for all future waterings.

There are two methods to add this solution. The first is to add the alcohol solution to what is already in the container. Add enough to bring it up to the proper level. The second will give shorter plants. In this second method, pour off all the old solution and replace it with the new each time additional solution is needed. So, how do we make the alcohol solution? An easy way is to use rubbing alcohol. Rubbing alcohol is usually 70% alcohol and should be mixed with 1-part alcohol with 10 or 11 parts water. Do not use beer or wine as the sugars present can interfere with normal growth. The researchers suggested the alcohol made it more difficult for the plants to take up water. This water stress stunted growth but did not affect the flowers

Starting Garden Transplants from Seed

January is often a cold and dreary month for many gardeners. However, planning for and starting vegetables and flower transplants from seed can make this a much more interesting time of year. Following are the steps needed for success in seed starting.





<u>Determine the date to seed:</u> There are two pieces of information that needs to be known in order to determine the date to seed transplants: the target date for transplanting outside and the number of weeks needed to grow the transplant. There is a companion article in this newsletter listing common plants and the number of weeks needed to grow a transplant. The target date for transplanting the cool-season crops such as broccoli, cabbage, cauliflower and onions are the end of March to the beginning of April. Warm-season crops like tomatoes, peppers and most annual flowers are usually planted about May 10 in Manhattan. Northern Kansas may be a week or so later than Manhattan and southern Kansas a week or so earlier.

seed remain viable for about 3 years, germination decreases as seed ages. See the accompanying article on using

Sowing Seed: Do not use garden soil to germinate seed as it is too heavy and may contain disease organisms. Use a media made especially for seed germination.

Keep Seed Moist: Seed must be kept moist in order to germinate. Water often enough that the media never dries. Using a clear plastic wrap over the top of the container can reduce the amount of watering needed. Remove the wrap after the seedlings emerge.

<u>Light:</u> Most plants will germinate in either darkness or light but some require darkness (Centurea, Larkspur, Pansy, Portulaca, Phlox and Verbena) and others require light (Ageratum, Browallia, Begonia, Coleus, Geranium, Impatiens, Lettuce, Nicotiana, Petunia and Snapdragon).

All plants require adequate amounts of light once emergence occurs. South facing windows may not provide adequate amounts and so fluorescent or LED fixtures are often used. Suspend fluorescent lights 2 to 4 inches above the top of the plants. LED lights are much more variable. Use LED lights that are designed to grow plants and follow the manufacturers recommendations. Regardless of the type of light used, leave the lights on for 16 hours each day.

<u>Temperature:</u> The temperature best for germination is often higher than what we may find in our homes especially since evaporating moisture can cool the germination media. Moving the container closer to the ceiling (top of a refrigerator) can help but a heating mat is best for consistent germination. A companion article lists common plants and their optimum germination temperature. After plants have germinated, they can be grown at a cooler temperature (65 to 70 degrees during the day and 55 to 60 degrees at night). This will help prevent tall, spindly transplants.

<u>Plant Movement:</u> Plants react to movement. Brushing over the plants with your hand stimulates them to become stockier and less leggy. Try 20 brushing strokes per day. However, brushing will not compensate for lack of light or over-crowding. Plants grown under inadequate light will be spindly regardless of any other treatment.

<u>Hardening Transplants:</u> Plants grown inside will often undergo transplant shock if not hardened off. Plants are hardened off by moving them outside and exposing them to sun and wind before transplanting occurs. Start about two weeks before transplanting and gradually expose the plants to outside conditions. Increase the number of hours and degree of exposure over the two-week period.

Larry E. Crouse
Butler County Horticulture Agent
K-State Research & Extension
206 N Griffith, Suite A
El Dorado, KS 67042
(316)321-9660
Icrouse@ksu.edu

old garden seed for more detailed information.

