

Kansas State University Cooperative Extension Office 206 N. Griffith, Suite A El Dorado, Kansas 67042



THE GRAPEVINE

February 13th, 2020

Larry Crouse Extension Horticulture Agent

HANDLING TIPS FOR VALENTINE'S DAY ROSES

If you are fortunate enough to receive roses from a loved one this Valentine's Day, follow these guidelines to help extend the life of your flowers.

For floral arrangements:

1) Keep the vase filled or floral foam soaked with warm water. Add fresh, warm water daily. If the water turns cloudy, replace it immediately. If possible, recut stems by removing one to two inches with a sharp knife. Do this under water. This allows the stems to draw in water, not air.

2) Keep flowers in a cool spot (65 to 72 degrees Fahrenheit), away from direct sunlight, heating or cooling vents, directly under ceiling fans, or near radiators.

3) If a rose starts to wilt, remove it from the arrangement, and recut the stem under water. Submerge the entire rose in warm water. The rose should revive in one to two hours.

For loose stems:

1) If you can't get your flowers in a flower food solution right away, keep in a cool place.

2) Fill a clean, deep vase with water and add the flower food obtained from your florist. Be sure to follow the directions on the package.

3) Remove leaves below the waterline. Leaves in water will promote bacterial growth.

4) Recut stems under water with a sharp knife and place the flowers in the vase solution.

Pruning Fruit Trees

Fruit trees can be pruned from now through March as long as the wood isn't frozen. Following are some general recommendations on pruning mature fruit trees followed by more specific instructions on each species.

General Recommendations

- Take out broken, damaged or diseased branches

- If two branches form a narrow angle, prune one out. Narrow angles are weak angles and tend to break during wind or ice storms.

- Take out all suckers. Suckers are branches that grow straight up. They may originate from the trunk or from major branches

- If two branches cross and rub against one another, one should be taken out.

- Cut back or remove branches that are so low they interfere with harvest or pruning. If cutting back a branch, always cut back to another branch or a bud. Do not leave a stub.

- Cut back branches to reduce the total size of the tree, if necessary.

- Thin branches on the interior of the tree.

Specific Instructions

<u>Peach and Nectarine:</u> Peach and nectarine require more pruning than any other fruit trees because they bear fruit on growth from the previous year. Not pruning results in fruit being borne further and further from the center of the tree allowing a heavy fruit crop to break major branches due to the weight of the fruit. Prune long branches back to a shorter side branch

<u>Apple:</u> Apples tend to become overgrown if not pruned regularly.

Wind storms and ice storms are then more likely to cause damage. Also, trees that are not pruned often become biennial bearers. In other words, they bear a huge crop one year and none the next. Biennial bearing

is caused by too many fruit on the tree. Though pruning helps, fruit often needs to be thinned as well. The goal is an apple about every 6 inches.

Spacing can vary as long as the average is about every 6 inches.

Cherry, Pear, Plum: Light pruning is usually all that is needed.

Simply remove branches that are causing or will cause a problem according to the general recommendations above.

Pruning Young Fruit Trees

Young fruit trees should be pruned to begin developing a strong structure of the main or scaffold limbs. This will help prevent limb breakage over the years when the scaffolds carry a heavy fruit load. Do not prune the year of planting except to cut back to the tree to a height of 36 inches. This will cause side branches to develop. If there are already side branches, do not prune the year of planting so the tree has the maximum number of leaves to help with establishment.

Apple, apricot, cherry, plum and pear trees generally are trained using the central leader system. The growth pattern for these trees is for a center branch to be dominant and to grow straight up. Peach and nectarine trees are normally pruned using the open center method because they do not have a strong tendency for one shoot or branch to dominate the growth of other shoots or branches. In this system, the tree is pruned to a vase-like pattern with no central leader.

Regardless of the system used, the three to four scaffold branches should:

o Be no lower than 18 inches from the ground. This makes it easier to prune and harvest.

o Form wide angles (about 60 to 80 degrees) with the trunk. Wide angles are much stronger than narrow angles and are less likely to break under wind or ice loads.

o Be distributed on different sides of the tree for good balance.

o Be spaced 6 to 10 inches apart on the trunk with none directly opposite or below another.

Check Plants for Scale Insects

The dormant season is a good time to check woody plants for scale insect infestations. This time of year, deciduous plants do not have leaves, so scale is more easily seen.

If an infestation is detected, make plans to apply a dormant oil for control by March 1. Be sure the temperature is 40 degrees or above before spraying. Scale insects are easily overlooked because they are small and immobile most of their lives and they do not resemble most other insects. Many of them resemble small shells that are oval or circular, but some have more unusual shapes like oyster shells. Coloring varies, but can include white, tan, and brown.

Plants that should be inspected for scales include apples, pears, other fruit trees, bush fruits, lilac, crabapple, oak, ash, elm, lilac, maple, linden, arborvitae, juniper, pine, spruce and yew. Manhattan euonymus, a broadleaf evergreen, is especially noted for having scale problems.

Plants are not harmed if only a few scales are present. But scale population can increase dramatically during the growing season. Heavy scale infestations can damage fruit crops, destroy branches and kill entire plants.

Larry E. Crouse Butler County Horticulture Agent K-State Research & Extension 206 N Griffith, Suite A El Dorado, KS 67042 (316)321-9660 Icrouse@ksu.edu

Kansas State University is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision, or hearing disability, contact Charlene Miller, KSRE - Butler Co. Director, 316-321-9660