

THE GRAPEVINE

April 30th, 2020

Larry Crouse

Extension Horticulture Agent

BAGWORMS, IT'S STILL TOO EARLY TO SPRAY

Timing is critical in many things, including controlling bagworms. Though handpicking is effective through much of the year, often it is impractical because of the sheer number of bagworms. However, if you only see a few bags, now would be a good time to pick them off and destroy them. As mentioned above, large populations of bagworms can make handpicking impractical. In such cases, spraying is recommended.

Bagworms only have one generation a year, and this makes controlling them a lot easier for us. Their eggs will be completely hatched and out of the bag by June 10th, so we target our sprays for the last three weeks of June and even into the first week of July.

Along with timing, thorough coverage is the other key to successful bagworm control. You must penetrate the canopy of the trees and shrubs you are spraying in order to impact the entire infestation. This requires a fairly powerful spray for evergreen trees and shrubs such as Red Cedars and other junipers.

As far as what to spray, this is the part of the equation that is of least importance for good bagworm control. Anything that is labeled for use as bagworm control will do a good job. Our top recommendation these days is 'Spinosad', a relatively safe and effective control that stops bagworms in their tracks. One of the formulations of Spinosad that can be found on the shelves of local nurseries and garden centers is 'Captain Jack's Dead Bug Brew'.

So there it is, but please wait until June before you spray or you will probably need to re-apply later in the summer.

Poison Ivy Identification and Control


Learning to identify poison ivy is vital if you wish to avoid the rash that accompanies exposure. Unfortunately, poison ivy can make identification difficult because it occurs in three forms: an erect woody shrub, a groundcover that creeps along the ground, and a woody vine that will climb trees. When poison ivy climbs, it forms numerous aerial roots that give the vine the appearance of a fuzzy rope. The leaves of poison ivy also vary. Though the compound leaf always has three leaflets, the leaf margins may be toothed, incised, lobed or smooth. The size of the leaves also can vary, although usually the middle leaflet is larger than the other two. Also, the middle leaflet is the only one with a long stalk; the other two are closely attached to the petiole (leaf stem). The number of leaves gives rise to the saying: "Leaves of three, let it be!" Poison ivy is often confused with Virginia creeper or Woodbine. Each of these vines, however, has five leaflets rather than three. There are three methods commonly used to eradicate poison ivy. These include pulling or grubbing out the plants by hand, cutting off the vine, and then treating the cut stump or the regrowth, and spraying the plants directly. The method used depends somewhat on the plant's growth form and how sensitive a person is to handling it. If the plant is growing as a groundcover, direct spray or grubbing the plant out is often used. If grubbing, wear gloves and a long-sleeved shirt. The soil must be moist for grubbing to work well. Wash the clothes and yourself immediately after you finish. It might also be a good idea to rinse the washing machine.


If the plant is in the shrub form, direct spray is the most common control method. If the plant is a woody vine that has climbed a tree, the preferred method is to cut the plant off at the base and treat the sprouts after they emerge. You don't need to worry about what is already up in the tree, once you separate the root system from the top, the top will die. Some triclopyr herbicides also have instructions on treating a freshly cut stump directly. Triclopyr (Brush-B-Gon Poison Ivy Killer, Hi Yield Brush Killer Stump Killer) is often used for poison ivy control. Other Herbicides that can be used include glyphosate (Roundup; Killzall Weed and Grass Killer; Nutgrass, Poison Ivy and Vine Killer). Poison ivy is tough. Repeat applications may be necessary.

Delay Planting Winter Squash and Pumpkins

Though early May is a good time to plant most of our warm-season crops such as tomatoes and peppers, it is best to delay planting of winter squash and pumpkins until about mid-June. Planting those two crops now will result in a crop that is mature enough to be attractive to the first generation of squash bugs in July. Delaying until about June 20 will allow plants to escape attack by the first generation. This later planting will also mean that pumpkins will mature closer to Halloween so that long-term storage is not needed. There will be a second generation of squash bugs that will hatch in August. Plant will need to be protected from that generation. See our publication on squash bugs for information on means of control.

Blackspot of Roses





A common disease of roses is blackspot, a fungus disease that can cause defoliation of susceptible plants. Look for dark, circular lesions with feathery edges on the top surface of the leaves and raised purple spots on young canes. Infected leaves will often yellow between spots and eventually drop. The infection usually starts on the lower leaves and works its way up the plant. Blackspot is most severe under conditions of high relative humidity (>85%), warm temperatures (75 to 85 degrees F) and six or more hours of leaf wetness. Newly expanding leaves are most vulnerable to infection. The fungus can survive on fallen leaves or canes and is disseminated primarily by splashing water. Cultural practices are the first line of defense.

1. Don't plant susceptible roses unless you are willing to use fungicide sprays. Purdue University has a list of blackspot resistant cultivars.
2. Keep irrigation water off the foliage. Drip irrigation works well with roses.
3. Plant roses in sun in areas with good air movement to limit the amount of time the foliage is wet.
4. Remove diseased leaves that have fallen and prune out infected rose canes.

If needed, protect foliage with a regular spray program (10- to 14-day schedule) of effective fungicides. Recommended fungicides include tebuconazole (BioAdvanced Disease Control for Roses, Flowers and Shrubs, BioAdvanced All-In-One Rose & Flower Care), myclobutanil (Immunox, F-Stop Lawn & Garden Fungicide) and chlorothalonil (Broad Spectrum Lawn & Garden Fungicide, Garden Disease Control, others).

Sweet Corn Primer

It used to be simple to decide which sweet corn to plant. You simply chose a cultivar and planted when the soil temperature reached 55 degrees. Now it has become more complicated due to genetic advances in sweet corn. Breeders have found certain genes that improve "standard" sweet corn. Below is an overview of the types commonly available to homeowners.

Standard (su): This is our "old" sweet corn and contains a "sugary gene" (su). Standard sweet corn should be isolated from field corn, popcorn, supersweets and ornamental corn. To isolate one type of corn from another, do not plant one type within 200 to 250 feet or be sure to have a difference of 12 to 14 days in time to maturity. Plant when the soil temperature reaches at least 55 degrees. Recommended varieties include Honey and Cream, Silver Queen, Sterling Silver, Jubilee, or Merit.

Supersweet (sh2): Though supersweets have up to three times the sweetness of standard sweet corns and hold their sweetness longer after harvest due to the sh2 gene, they do have some drawbacks such as tougher kernels and a lack of some of that good "corn" flavor. They also need to be isolated from other sweet corn types and are very sensitive to cooler soils. Wait until the soil temperature reaches 65 degrees before planting. Try Candy Store, Florida Staysweet, Sugar Loaf, Sweet Time, or Sweetie.

Sugar Enhanced (se): These are probably the most popular type of sweet corn grown due to their tender kernels, good flavor and less sensitivity to cool soils (60 degree soil temperature for planting). They hold their post-harvest sweetness longer than standard types but will not hold sweetness as long as the supersweets. The sweetness from the sugar-enhanced types is due to the "se gene." If both parents were se types, the variety is known as an se+ or se se. If only one parent was an se type and the other an su type, then the variety will be listed as se. They do not need to be isolated other than from the supersweets. Suggested varieties include Bodacious, Ambrosia, Sweet Temptation, Delectable and Miracle.

Triplesweet (synergistic): The newest types of sweet corns blend the su, se and supersweet types with the goal of combining the best characteristics of each. We don't have firm recommendations yet but you may want to try Serendipity, Polka, Avalon or Frisky.

Larry E. Crouse
Butler County Horticulture Agent
K-State Research & Extension
206 N Griffith, Suite A
El Dorado, KS 67042
(316)321-9660
lcrouse@ksu.edu

Kansas State University is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision, or hearing disability, contact Charlene Miller, KSRE - Butler Co. Director, 316-321-9660

